# Span

SHELL PUBLIC HEALTH AND AGRICULTURAL NEWS

# Index

VOLUME 1. NUMBERS 1-4 APRIL 1958 TO JANUARY 1959

Digitized by the Internet Archive
in 2025

https://archive.org/details/span-1958_april-1958-january-1959_1_index

# Span

VOLUME 1 - NUMBERS 1-4

## AUTHOR INDEX


Allan, P. . . . . (2) 2
Busvine, J. R. . . . (1) 8, (2) 8
Caseby, W. R. . . . . (2) 24
Comins, P. C. M. . . . . (1) 14
Courshee, R. J. . . . . (4) 13
Crafts, A. S. . . . . (3) 5
Davidson, G. . . . . (3) 12
Fenwick, D. W.. . . . . (2) 5
Grist, D. H. . . . . (4) 27
Hamman, R. E.. . . . . (4) 34
Harris, A. B. . . . . (3) 43
Harrison, J. W. . . . . (3) 30
Little, E. C. S. . . . . (1) 25
Meijneke, C. A. R. . . . . (4) 2
Metcalfe, C. B. . . . . (3) 34
Oren, R. . . . . (1) 17
Parkin, Anthony . . . . (1) 4
Pearson, A. J. A. . . . . (3) 2
Reid, T. N. . . . . (1) 30
Richardson, A. G. . . . . (2) 28
Sandford, H. . . . . (1) 21
Sanger, A. M. H. . . . . (3) 16
Skinner, J. E. . . . . (4) 23
Stanford, C. J. N. . . . . (4) 20
Wain, R. L. . . . . (4) 9
Webster, B. N. . . . . (2) 34
Wilkins, V. E. . . . . (1) 20
Wilson-Jones, K. . . . . (3) 26
Wood, P. L. . . . . (4) 6
Wright, P. . . . . (4) 31

# SUBJECT INDEX

## A


Acaricide, Phosdrin as, (2) 29, 30
Accidents, farm, analysis of causes, (1) 28
  overturning tractors, (1) 28
*Aedes aegypti*, generations per year, (2) 11
  insectide resistance, (1) 11; (2) 10
*Aedes dorsalis*, (1) 11
*Aedes nigromaculis*, (1) 11
*Aedes sollicitans*, (1) 11
Aerial spraying. *See* Spraying.
*Agave sisalana*. *See* Sisal.
Agricultural Advisory Service, Rhodesia, (3) 38
Agricultural chemicals, (1) 2
  hazards from use of, (1) 14
  Shell's research work, (3) 23
Agricultural Research Centre, Woodstock, (3) 22
Agricultural Research Laboratory, Modesto, (3) 23
Agricultural worker, instruction in tractor driving, (1) 28
Agriculture (Poisonous Substances) Regulations (1) 14
Aircraft, use in agriculture, (1) 25; (2) 24; (3) 33
Aldrin, (3) 13
  effect on beneficial insects, (3) 16, 18, 19
  pest resurgence after use, (3) 19
  resistance build-up in housefly, (2) 10
  timing, on melon, (3) 18
  toxicity to ectoparasites of sheep, (4) 32
    to sisal weevil larvae, (1) 23
    to tsetse fly, (4) 24
  use on sisal plantations, (1) 21, 24
Aldrin/diatomite dust, (1) 23
Allyl alcohol, toxicity to broomrape, (2) 14
Aminotriazole, movement in plant tissues, (3) 6
  selective weed killer, (4) 10
Ammonium nitrate, foliar application, (2) 2, 3
Ammonium sulphate, for sisal, (1) 22
*Amorbia emigratella*, resurgence, (3) 18
Analysis of pesticide residues, (2)33, BR*; (3) 25
*Anopheles*, generations per year, (2) 11
*Anopheles atroparvus*, (1) 9
*Anopheles culicifacies*, (1) 9; (3) 14
*Anopheles darlingi*, (1) 9; (2) 9
*Anopheles gambiae*, (2)31 BR; (3) 14
  generations per year, (2) 11
  insecticide resistance build-up, (2) 10
  resistance to dieldrin, (1) 10; (2) 8
    to dieldrin-BHC, (3) 12
*Anopheles labranchiae*, (2) 11
*Anopheles maculatus*, (2) 9; (3) 14
*Anopheles pseudopunctipennis*, (1) 9
*Anopheles quadrimaculatus*, (1) 10
  resistance to dieldrin-BHC, (3) 12
*Anopheles sacharovi*,
  resistance to DDT, (1) 9; (3) 12
  resistance to dieldrin-BHC, (3) 13
*Anopheles stephensi*, (1) 10 (fig.)
  control by dieldrin, Saudi Arabia, (4) 18
  resistance to DDT, (1) 10; (3) 12
*Anopheles subpictus*, (3) 13
*Anopheles sundaicus*, (2) 31 BR
  resistance to DDT, (1) 10; (3) 12, 13
*Anopheles taeniorhynchus*, (1) 11
*Anthocoris nemorum*, (3) 17 (fig.)
Antimalarial drugs, (2) 31 BR
*Apanteles glomeratus*, (3) 17 (fig.)
*Aphelenchoides cocophilus*, (2) 5, 6
Aphids, control by Phosdrin, (2) 29
  destroyed by lady bird larvae, (3) 19
  East African species, (4) 38 BR
Apple, aerial spraying trial, (2) 27
  copper deficiency symptoms, (4) 7 (fig.)
  foliar absorption of nutrients, (2) 3
  Mediterranean fruit fly on, (1) 17
  seedling, nematode damage to root, (4) 3 (fig.)
  pre-harvest drop, control, (4) 10
  zinc deficiency symptoms, (4) 7 (fig.)
Apple stocks, protection by D-D, (4) 5 (fig.)
Apricot, Mediterranean fruit fly on, (1) 17, 19
Aquatic weed, control, (3) 26
Army worm, control by dieldrin, (1) 27
Asparagus, tolerance of urea spray, (2) 3
*Asystasia gangetica*, control, (1) 24
Atomic energy, role in agriculture, (1) 6
Atomizer, for aerial spraying, (2) 25




* Book Review

Auster Agricola aircraft, (2) 26 (fig.)
Auster Aiglet, (2) 26
Auster Workmaster, (2) 24, 27, (fig.)
Australia, fertiliser usage, (3) 41 BR
sheep ectoparasites, (4) 31
wool industry losses through pests, (4) 32
Autoradiographs, barley plants, (3) 8–11 (figs.)
Autoradiography, systemic insecticides (3) 7
weedkiller penetration of tissues, (3) 6, 8–11 (fig.)

## B

Ballasting of tractors, (2) 29
Banana, foliar absorption of urea, (2) 4
leaf spot, control by mineral oil, (1) 3; (3) 2, 3 (fig.)
Barley, mildew-resistant, radiation mutant (1) 4
translocation of weedkiller in, (3) 7, 8–11 (figs.)
Bawden, F.C. biographical note, (3) 15
"Bayer 21/199" resistance of flies to, (2) 11
Beans, tolerance of foliar sprays, (2) 3
Bed bug, control by DDT, (1) 12 (and fig.)
insecticide resistance build-up, (1) 12
Beef production, cattle breeds for tropics, (4) 21
Belgium, fertiliser usage, (3) 41 BR
Benochlor, (3) 28
Benzene hexachloride, (3) 13
resistance to, mechanism, (2) 10
of anopheline mosquitoes, (1) 10; (3) 12
of *Culex fatigans*, (1) 12
toxicity to body louse, (1) 12
to coccinellids, (3) 18
to ectoparasites of sheep, (4) 32
to mosquitos, (1) 10, 11; (2) 31, BR
to tsetse fly, (4) 24
Bindweed, movement of weedkillers in, (3) 5, 6
Bioassay, of pesticide residues, (2) 33 BR; (3) 24
Biological control, of insect pests (2) 33 BR
of pondweeds, by *Tilapia* sp, (3) 27
Biuret, limit for urea, (2) 2, 4
Black scale of citrus, predator of, (1) 20
*Blatella germanica*, control, (1) 13
generations per year, (2) 11
insecticide resistance, (1) 13; (2) 10
Blister blight of tea, (2) 34
correlation of incidence with sunshine hours, (2) 35
Blowfly, damage and control, (4) 31, 33 (fig.)
resistance to dieldrin reported, (4) 32
Body louse, insecticide resistance, (1) 12; (2) 9
parasite of sheep, (4) 32, 33 (fig.)
Book reviews,
Aphididae of East Africa, by Eastop, (4) 38
Beneficial insects, by Moreton, (2) 33
British crop husbandry, by Sanders, (4) 37
Comparison of high- and low-volume spraying techniques, by S.C.I. Pesticides Group, (2) 33
Cotton, by Brown and Ware, (3) 40
Farming weather, by Smith, (4) 38
Fertiliser progress, by F.A.O., (3) 41
Flora of the British Isles, by Clapham and Tutin, (3) 42
Insect migration, by Williams, (3) 39
Insecticide resistance, by Brown, (4) 40
Malaria control, by Macdonald, (2) 31
Marshes of South Iraq, (3) 43
Pest control research, by Metcalf, (2) 33
Soil: The 1957 Year book of Agriculture, Ed. by Stefferud, (2) 33
Tea, by Eden, (4) 36
Tropical Agriculture, by ICTA, (4) 40
Veterinary toxicology, by Garner, (4) 39
Weed control, by Helgeson, (3) 42
World of the Soil, by Russell, (2) 32
*Boophilus decoloratus*, (2) 10
Borax, use on root crops, (4) 7
Bordeaux mixture, compatible with urea in sprays, (2) 3
Boron deficiency, in crop nutrition, (4) 6, 7
Boron toxicity, correction by drainage, (4) 7
Braconid fly, (3) 17 (fig.)
Bradford Cattle, (4) 21
Brahman cattle, cross with beef breeds, (4) 21
Brangus Cattle, (4) 21
Breech strike of sheep, (4) 31
control by Mulesing, (4) 32
Breeding, plant. *See* Plant breeding,
Breeding Stock, degeneration, (4) 22
export from Great Britain, (4) 21
British Weed Control Conference, (4) 26
Broom, eradication by 2, 4, 5-T, (1) 26
Broomrape, (2) 15 (fig.)
control by allyl alcohol, (2) 14
Brown heart, of swedes, control, (4) 7
Brush control, (4) 26
Buckrake, (2) 18 (fig.) 19
Buffalo bean, control in sisal field, (1) 24
'Burgo', (3) 33
Bush clearance, in tsetse fly control, (4) 25
Butyric herbicides, selectivity of, (4) 12

## C

Cabbage, tolerance of foliar spray, (2) 3
Cabbage looper, control by Phosdrin, (2) 30
Calcium cyanide residues, zero tolerance, (4) 35
Canada, fertiliser usage, (3) 41 BR
Canadian pondweed, spread in waterways, (3) 29

Canal, weed clearance, (3) 27, 28 (fig.)
Captan, (1) 3
Carbon-14, herbicide translocation study, (3) 5
Carrot, tolerance to urea sprays, (2) 3
Cat flea, resistance to DDT, (1) 12
Caterpillar, aerial spraying, New Zealand, (1) 27
Cattle, British Breeds, (4) 20 (fig.) 21
  nagana incidence, Africa, (4) 23
  population, New Zealand, (1) 25
Cattle baits, in tsetse fly control, (4) 24
Cauliflower, tolerance of foliar spray, (2) 3
*Ceratitis capitata*, *See* Mediterranean fruit fly.
*Ceratophyllum*, control, (3) 27
*Cercospora musae*, (1) 3; (3) 2, 4
Cereals, copper deficiency symptoms, (4) 8
  production, Turkey, (1) 32
  radiation mutants, (1) 4
  tolerance of nitrogen sprays, (2) 2
  weed control in, (4) 10, 12
Ceylon, blister blight of tea, (2) 34
  tea production, (4) 36 BR
*Chara*, eradication by dredging, (3) 27
Chelating compounds, (4) 8
Cherry, aerial spraying trial, (2) 27
*Chilocorus bipustulatus*, (1) 20
Chlordane, (3) 13
  resistance to, mechanism, (2) 10
    of anopheline mosquitoes, (1) 9, 10
    of cockroach, (1) 13
  toxicity to cat flea, (1) 12
Chlorinated hydrocarbon insecticides, formulation, (3) 31
  resistance to, of disease vectors, (1) 8; (2) 8
    *See also* Aldrin, Benzene hexachloride, Chlordane, Dieldrin, Endrin, Heptochlor, Isodrin.
Chloropicrin, fumigant efficiency, (4) 4
Chloropropham, (4) 10
*Cimex hemipterus*, (1) 12
*Cimex lectularius*, (1) 12 (fig.)
Citrus, copper deficiency, (4) 6, 7
  fumigation of fruits with ethylene dibromide, (2) 36
  Mediterranean fruit fly on, (1) 18
  virus diseases, EPPO project, (2) 23
Citrus Marketing Board, (1) 18, 20
Clamp, silage, (2) 18 (fig.)
Cleavers, control by MCPP, (4) 12, 26
Climatology, micro-, (4) 38, BR
CMPP, toxicity to cleavers, (4) 12, 26
Coccinellids, reaction to insecticides, (3) 18
*Coccus hesperidum*, resurgence, (3) 18, 19
Cockroach, insecticide resistance, (1) 13
Coconut, red ring disease, (2) 5
Coconut weevil, damage by, (2) 6
Codling moth, control, and orchard pest resurgence, (3) 20
Coffee, foliar absorption of urea, (2) 4
Colorado beetle, international control, (2) 21, 23 (fig.)
*Commelina benghalensis*, (1) 24
*Compsiluria concinnata*, (3) 17 (fig.)
Conservation, soil and water, Rhodesia, (3) 34
*Convolvulus arvensis*, (3) 5, 6
Copper deficiency, apple tree, (4) 7 (fig.)
Copper fungicide, against blister blight, (2) 35
Copper-in-oil concentrate, aerial spray, (2) 27
Copper salts as selective weedkillers, (4) 10
Copper sulphate, against water weeds, (3) 28
Cost, aerial spraying, (1) 26
  desert locust control, (3) 44
  malaria eradication, Mexico and India, (2) 4
  Mediterranean fruit fly campaign, Florida, (2) 36
  nematode control in tree nurseries, (4) 4
*Costelytra zealandica*, (1) 27
Cotton, acreage, Turkey, (1) 32
  diseases of, (3) 40 BR
  production, (3) 40 BR
Crategus, growth habit on exhausted soil, (4) 2
*Crategus oxycantha*, (4) 4
Crawler tractor, stability, (2) 30
Crop husbandry, (4) 37 BR
  in tropics, (4) 40 BR
Crop protection, international action, (2) 20
Crops, pesticide residue tolerances, (4) 34
  spray drift hazard to, (4) 16
  statistics, Turkey, (1) 32
Cryolite, insecticidal value, (1) 20
*Ctenocephalides felis*, (1) 12
Cuba, tobacco pest, (2) 14
Cucumber, sensitivity to urea spray, (2) 3
Cucurbita, sensitivity to foliar sprays, (2) 2, 3
*Culex fatigans*, (1) 11; (2) 9, 10
*Culex lectularius*, (2) 10
  generations per year, (2) 11
*Culex tarsalis*, (1) 11; (2) 10
Cuttings, rooting stimulants, (4) 10
*Cystus* sp., (1) 26

## D

2, 4-D, (4) 10
  aerial application, (1) 28
  control of aquatic weeds by, (3) 28, 28 (fig.)
    of water hyacinth, (3) 29
  movement in plant, tracer study, (3) 6, 8–11 (fig.)
  susceptibility of willow to, (1) 26
  use in sisal, (1) 24

Dairy cow, silage in rations of, (2) 16, 17 (fig.)
Dalapon, (1) 3; (4) 10, 26
movement in plant tissues, (3) 7
use in sisal nursery, (1) 24
Dalgrond, definition, (4) 5
*Damalina ovis*, (4) 32, 33 (fig.)
DANP, resistance of body louse to, (2) 9
2,4-DB, selective weedkiller, (4) 12
D-D, application method, (4) 4
comparative fumigant efficiency, (4) 4
persistence in soil, indicator crop, (4) 5
DDT, content in normal diet, (1) 15
mosquito control, Mexico, (4) 17
resistance to, (1) 3; (4) 40 BR
first report of, (2) 12
genetical aspect, (2) 9, 11; (3) 12
mechanism in house fly, (2) 9
of body louse, (1) 12; (2) 9
of disease vectors, (1) 8; (2) 8
of fleas, (1) 12
of mosquito species, (1) 9, 10, 11, (2) 31 BR; (4) 18
sprays for citrus, (1) 20
for oaks, (3) 33
toxicity to *Anopheles gambiae*, (1) 10
to coccinellids, (3) 18
to ectoparasites of sheep, (4) 32
to grass grub, (1) 27
to ladybird predators, (1) 20
to tsetse fly, (4) 24
DDT-dehydrochlorinase enzyme, in fly, (2) 9
Deficiency diseases of plants, (4) 6
Defoliation, studies on cotton, (3) 40 BR
Demeton, (2) 30
Desert locust, (2) 35
control, economics, (3) 44
Dianisyl neopentane, insect resistance to, (2) 9
Diatomite dust, filler for aldrin, (1) 23
Diazinon, (2) 11, 31 BR
2,4-Dichlorophenoxy butyric acid. *See* 2, 4-DB.
Dichloropropane-dichloropropene. *See* D-D.
Dieback of citrus, (4) 7
Dieldrex 15, (1) 23
Dieldrin, (2) 31 BR; (3) 32
control of DDT resistant anophelines by, (3) 12
effect on beneficial insects, (3) 16, 18, 19
formulation, (2) 35; (3) 31
granular, uses, (2) 36
mosquito control, Mexico, (4) 17
Saudi Arabia, (4) 18
pest resurgence after use, (3) 18
resistance to, build-up in fly, (2) 10
of cockroach, (1) 13
of mosquitoes, (1) 9, 12; (2) 8; (3) 12
sprays for citrus, (1) 20
timing on melon, (3) 18
toxicity to *Aedes aegypti*, (1) 11
to *Anopheles stephensi*, (1) 11
to *Anopheles sundaicus*, (1) 10
to army worm, (1) 27
to ectoparasites of sheep, (4) 32
to sisal weevil larvae, (1) 23
to tsetse fly, (4) 24, 25
use in sisal plantations, (1) 21, 24
Dinitro compounds toxicology, (4) 39 BR
Dinitro-*ortho*-cresol. *See* DNOC.
Dinitro-*sec.*-butylphenol. See Dinoseb.
Dinoseb, as pre-emergence weed spray, (4) 10
Dithane, compatible with urea in spray, (2) 3
DNOC, (4) 10
fatalities on contractors' workers, (1) 14
residues, zero tolerance, (4) 35
Drainage systems, weed control in, (3) 26
Drift from ground sprayers, (4) 13
preventive means, (4) 14, 15 (fig.)
use of invert emulsions, (4) 26
Droplet size, choice of, (2) 25
in mineral oil sprays, (3) 3
relation to spray drift, (4) 15 (fig.)
spray efficiency and, (2) 25
*Drosophila melanogaster*, (2) 9
Dust concentrates, formulation, (3) 32
Dusting crops with dieldrin, (3) 32
Dusts, formulation, (3) 31

## E

East Africa, aphids found in, (4) 38 BR
Ectoparasites of sheep, Australia, (4) 31
Edgar Percival EP9, (2) 24, 26 (fig.)
*Eichornia crassipes*, (3) 29
*Elodea canadensis*, (3) 27, 29
*Empoasca fabae*, resurgence after sprays, (3) 18
Emulsifiable concentrate, (3) 31
Emulsions, breaking, control of, (3) 31
Endrin, (3) 13
effect on beneficial insects, (3) 16, 18, 19
granular formulation, (3) 32
resistance build-up, in housefly, (2) 10
selective toxicity, (3) 20
sprays for oaks, (3) 33
toxicity to mouse, (3) 18
Enzyme, dehydrochlorinating, in fly, (2) 9
Epidemics, flea-borne, control, (1) 12
louse-borne, control, (1) 12
EPPO (2) 20 (map), 21
Erosion, soil, by rainstorms, (3) 35
by water course, (3) 34 (fig.)

Ethylene dibromide, (4) 4
dip for citrus fruits, (1) 18
fumigation of fruits and vegetables, (2) 36
European and Mediterranean Plant, Protection Organisation (EPPO), (2) 20
European corn borer, control by endrin, (3) 32
*Euphorbia hirta*, (1) 24
Exanthema of citrus, (4) 7
Exhaust aerosol application, (2) 35; (4) 24

## F

Fall webworm, international action, (2) 22
Farm machinery, accidents, (2) 28
Farm planning service, Rhodesia, (3) 38
Farm worker, instruction in tractor driving, (1) 28
Feeding trials with irradiated foods, (1) 7 (fig.)
Ferbam, (2) 3
Fern, aquatic, control, (3) 26
Fertiliser research, tracer technique, (1) 5
Fertilisers, aerial application, (2) 26
foliar application, (2) 2
for fish pond plankton, (3) 27
for rice, economics of use, (4) 30
granular, application, (2) 27
statistics of production and use, (3) 41 BR
uses, (4) 37 BR
Fibre crop, yield from dieldrin spraying, (1) 24
from nitrogen fertiliser, (1) 22
Field strength dusts, formulation, (3) 32
Fig, Mediterranean fruit fly on, (1) 18
production, Turkey, (1) 32
Filariasis, diethylcarbamazine against, (4) 18
Fillers, for dust concentrates, (3) 32
Fire ant, control campaign, U.S.A., (2) 36
Fission product. *See* Radioactive isotope,
Flea, resistance to DDT, (1) 12
Florida, Mediterranean fruit fly, (2) 36
Flora, British, (3) 42 BR
Flowering habit, sisal, (1) 22
Foliar application of nutrients, (2) 2
ammonium nitrate to vegetables, (2) 2, 3
urea, (2) 2, 3
to tropical crops, (2) 4
Food and Agriculture Organisation, aid for fall webworm control, (2) 22
Food and Drug Administration, U.S.A., rules on pesticide residues, (1) 15
Food preservation, irradiation method, (1) 6
Forage harvester, (2) 18 (fig.) 19
Forest plantation, weed control in, (1) 26
Formulation, of agricultural chemicals, (3) 25
of pesticides, (3) 30
of seed disinfectants, (3) 32
France, fertiliser usage, (3) 41 BR
Freeze drying, of plants for tracer studies, (3) 6
Fruit trees, damage by Mediterranean fruit fly, (1) 17
spraying techniques, (2) 33 BR
zinc deficiency, (4) 7 (fig.) 8
Fumigants, soil, (1) 3
Phosdrin as, (2) 28
*see also* D-D, Ethylene dibromide, Methyl bromide, Vapam
Fumigation of tree nursery, cost, (4) 4
Fungicides, (1) 3; (2) 27
cost reduction, (2) 34
mineral oils, evaluation, (3) 2
screening tests, (3) 25
*see also* Captan, Ferbam
Fungitoxicity, mechanisms, (2) 33 BR

## G

*Galium aparine*, (4) 12, 26
Game animals, tsetse fly reservoir, (4) 25
Genetic factors in resistance build-up, (2) 9, 11, 31 BR
(3) 12
*Geocoris*, sensitivity to aldrin, (3) 18
Gibberellins, (1) 3
Glasshouse research, Modesto, (3) 23, 25 (fig.)
Glossary, weed control terms, (3) 42 BR
*Glossina brevipalpis*, (4) 24
*Glossina morsitans*, (4) 23, 25
*Glossina pallidipes*, (4) 24
*Glossina palpalis*, (4) 23, 25
Goat, population of Turkey, (1) 32
Gorse, eradication, (1) 25
use of 2,4,5-T, (1) 26
Granular formulation, (3) 31
Granules, insecticide, penetrative power, (3) 32
Grape production, Turkey, (1) 32
Grass, ensilage, (2) 16
harvesting machines, (2) 18, 19
Grass grub, control by DDT, (1) 27
Grass species, incompatibility, removed by irradiation, (1) 4
weedkillers for, (3) 26; (4) 10, 26
Grassland, foliar application of fertilisers (1) 3
productivity, U.K., (3) 21
soil conserving value, (3) 37
topdressing by aircraft, (1) 25
Great Britain, agricultural chemicals Notification and Clearance Scheme, (1) 16
Greece, antimalarial campaign, (1) 9
Green crop loader, (2) 18 (fig.) 19
Guava, Mediterranean fruit fly on, (1) 18

## H

Harvesting, machinery, (2) 18 (fig.)
of sisal, (1) 22
Hazel nut, production, Turkey, (1) 32
Health hazard, from pesticide residues in crops, (1) 14
Heart rot of sugar beet, (4) 7
Helicopter, spraying from, (1) 25, 26 (fig.)
of locusts, (2) 24
of tsetse fly, (4) 24
Heptachlor, granular, use, (2) 36
resistance mechanism in fly, (2) 10
Herbicides. *See* Weedkillers.
Hereford, cross with Brahman breed, (4) 21
Hexaethyl tetraphosphate, (4) 35
Highland cattle, in the Andes, (4) 21 (fig.)
*Homona coffearia*, (3) 18
Hormones. *See* Plant growth regulators; Weedkillers.
Host plants, of aphids, (4) 38 BR
Housefly, (1) 11 (fig.)
control by organo phosphorus compounds, (1) 9; (2) 29
disease vector, (1) 8
generations per year, (2) 11
resistance build-up, (1) 8; (2) 8
mechanism discussed, (2) 9, 10; (4) 40 BR
*Hyalopterus arundinis*, (3) 18
Hydrogen cyanide, residues, tolerances, (4) 35
resistance to, in scale insects, (2) 9
*Hyphantria cunea*, (2) 22

## I

IAA, movement in plant tissues, (3) 7
IDLIS (3) 44
India, fertilizer usage, (3) 42 BR
losses due to malaria, (2) 4
malaria eradication, cost of, (2) 4
Insect pests, biological control, (2) 33 BR
of stored products, (2) 23
resurgence after spraying, (1) 20; (3) 16, 18, 19
Insect repellents, (2) 33 BR
Insecticide, aerial application, (1) 27; (4) 24
droplet size, (2) 25
emulsifiable concentrate, (3) 31
formulation, (3) 30
residues, (1) 3, 15, 16 (fig.) (2) 33 BR; (4) 34
selective toxicity concept, (2) 11
systemic, (2) 28; (3) 20
vigour tolerance to, (2) 9; (3) 13
Insecticide resistance, (1) 2; (2) 31 BR; (4) 40 BR
blowfly to aldrin and dieldrin, (4) 32
build-up in disease vectors, (1) 8; (2) 8
counter measures, (2) 12
genetic aspects, (2) 9, 11, 31, BR; (3) 12
Insects, beneficial, (2) 33 BR; (3) 17 (figs.)
reduction by insecticides, (1) 20; (3) 16
disease-carrying, (1) 8; (2) 8
seasonal migration, (3) 39 BR
Instability of tractors, (1) 28
International Colorado Beetle Committee, (2) 23
International Congress on Tropical Medicine and Malaria, Sixth, (4) 17
International Co-operation in crop protection, (2) 20 (map), 21
International Desert Locust Information Service (IDLIS), (3) 44
Iraq, antimalarial campaign, (3) 43 BR
Iridiomyrmex, in guava, killed by dieldrin, (3) 19
Iron deficiency, in plant nutrition (4) 6, 8
Irradiation laboratory, of A.E.R.E., (1) 4
Irrigation systems, weed control in, (3) 26, 29 (fig.)
Isodrin, (2) 10; (3) 13
Itch mite, of sheep, (4) 32, 33 (fig.)

## J

Jamaica, banana leaf spot control, (3) 2
Jamaica Red cattle, (4) 21
Japan, fertiliser usage, (3) 41 BR
rice growing methods, (4) 28
Java, antimalarial campaign, (1) 10
Jersey cattle breed, in Kenya, (4) 20 (fig.)

## K

Ked, sheep, control, (4) 32, 33 (fig.)
Kenya, sisal growing, (1) 21
tsetse fly campaign, (4) 25
Knapsack sprayer, double nozzled, (2) 34
viscosity of mineral oils for, (3) 4
Koela forage harvester, (2) 18 (fig.)

## L

Labelling agricultural chemicals, (1) 14, 15 (fig.)
*Laburnum anagyroides*, (4) 4
Ladybirds, beneficial, (1) 20; (3) 19 (fig.)
Lake weeds, control, (3) 26
Land use, Turkey, (1) 31

Large Milkweed Bug, (2) 29
Larvicides, (2) 12
Phosdrin as, (2) 30
Lead arsenate, (2) 36
Leaf scorch, of banana, (3) 2
Legislation, agricultural chemicals, (1) 3, 14, 15
movement of sheep, Australia, (4) 32
Natural Resources Act, Rhodesia, (3) 36
pesticide residues, U.S.A., (4) 34
plant protection, (2) 22
Legumes, weed control in, (4) 12
*Leptinotarsa decemlineata*, (2) 21, 23 (fig.)
*Leptospermum* spp., (1) 26
Lettuce, indicator crop, D-D persistence, (4) 5
Lettuce aphid, control by Phosdrin, (2) 29
Liming, of sisal, (1) 22
trace element availability and, (4) 8
*Liriomyza subpusilla*, (3) 18
Lisbon Conference, (4) 17
Livestock, export from Great Britain, (4) 20, 21
statistics, New Zealand, (1) 25
Turkey, (1) 32
Locust, breeding cages, (3) 24 (fig.)
control, exhaust-nozzle spraying, (2) 35
information service, (3) 44
migration, (3) 39 BR
Low-volume spraying, (2) 33 BR
droplet size, (2) 25
Sayer exhaust nozzle for, (2) 35
Lucerne, foliar absorption of potash, (2) 2
spotted aphis, sensitive to Phosdrin, (2) 29
*Lucilia cuprina*, (4) 31
*Lucilia sericata*, (4) 31

## M

*Macrosiphum pisi*, (2) 29
resurgence after spraying, (3) 18
Maize, as soil-conserving crop, (3) 37
Malaria, epidemology and control, (2) 31 BR
eradication, Lisbon Conference, (4) 17
South Iraq methods, (3) 43 BR
WHO programme, (1) 27; (3) 44
funds of WHO and PASO, (1) 27
national campaigns, Greece, (1) 9
Java, (1) 10
Nigeria, (1) 10
Trinidad, (1) 11
WHO survey, (2) 4
Malariolists at Lisbon, (4) 17
Malathion, (3) 13
aerial sprays against Mediterranean fruit fly, (2) 36
cockroach control by, (1) 3
comparative efficiency, (2) 29, 30
mosquito susceptibility to, (2) 31 BR
Maleic hydrazide, movement in plants, (3) 6, 10–11 (figs.)
*Malus pumilia*, (4) 4
Manganese sulphate, use, (4) 8
Manganese toxicity, to kale, (4) 7 (fig.)
Mango, fumigation with ethylene dibromide, (2) 36
Manuka, eradication by 2,4,5-T, (1) 26
*Marmara* sp., resurgence, (3) 18
Marrow stem kale, manganese toxicity, (4) 7 (fig.)
McIlwaine commission, work of, (3) 35
MCPA, (1) 25; (4) 10
aquatic weed control by, (3) 28
mixture with 236-TBA, (4) 26
use in sisal, (1) 24
MCPB, selective weedkiller, (4) 11 (fig.) 12
MCPP, selective weedkiller, (4) 12
Mealy bug, ant attendants killed by dieldrin, (3) 20
Meat preservation, by radiation, (1) 6
Mechanisation, of rice culture, (4) 28
of silage crop harvesting, (2) 18, 19
Mediterranean fruit fly, (1) 17 (fig.)
control, economics of, (1) 19
eradication, Florida, (2) 36
EPPO report, (2) 23
life cycle, (1) 18
pupa, (1) 18, (fig.)
quarantine regulations, (1) 18
spray programme for citrus, (1) 20
*Melophagus ovinus*, (4) 32, 33 (fig.)
Mercurised copper oxychloride, aerial application to potatoes, (2) 27
Mercury-containing compounds, residues, zero tolerances, (4) 35
Meteorology for farmers, (4) 38 BR
Methyl bromide, fumigation of vegetables, (2) 36
Methyl-4-chlorophenoxyacetic acid. *See* MCPA.
Methyl-4-chlorophenoxybutyric acid. *See* MCPB.
Methyl-4-chlorophenoxypropionic acid. *See* MCPP.
Methoxone, against aquatic weeds, (3) 28
Methoxychlor, sprays for citrus, (1) 20
toxicity to coccinellids, (3) 18
Mexican bean beetle larvae, (2) 29
Mexico, malaria eradication, (2) 4
Micro-bioassay, of residues, (2) 33; (3) 24 (fig.)
Micronair rotary atomiser, (2) 25, 27 (fig.)
Migration of insects, (3) 39 BR
Mildew resistance, breeding for, (1) 4
Miller Amendment, (1) 15; (4) 34
Mineral oils, agricultural uses, (3) 23
in banana leaf spot control, (1) 3; (3) 2, 4
mosquito larvicides, (3) 23
with pentachlorophenol, as weedkiller, (3) 29

Mist blower (Kiekens-Dekker), (4) 25
Mite, resurgence after spraying, (3) 18
sensitivity to Phosdrin, (2) 29, 30
Modesto Agricultural Laboratory, (3) 23
Molybdenosis, in ruminants, (4) 8
Molybdenum deficiency, in plants, (4) 8
Monuron, (4) 10
movement in plant tissues, (3) 7
Mosquito, control, (1) 27; (2) 31 BR; (4) 17
insecticide resistance, (1) 9, 11; (2) 9, 31 BR; (3) 12
test kit devised by WHO, (2) 8, 13 (fig.)
Moth-proofing emulsions, formulation, (3) 32
Mouse, control in orchards, (3) 18
*Mucuna pruriens*, control, (1) 24
Mulesing, of sheep, (4) 32
*Musca domestica*. *See* Housefly.
Mustard, mildew-resistant mutant, (1) 4
Mutation, induced, in rice, (4) 30
radiation-induced, (1) 4, 5
*Mycosphaerella musicola*, (3) 2

## N

*Nabis*, sensitivity to aldrin, (3) 18
Nagana, in cattle, (4) 23, 25 (fig.)
*Najas*, floating weed, control, (3) 27, 28
Naphthenic oils, (3) 4
phytotoxicity, (3) 2
Nemagon, (2) 7; (4) 4
Nematocides, *See* Chloropicrin, D-D, EDB, Nemagon, Vapam.
Nematode, count, (4) 3
disease of coconut, (2) 5
soil exhaustion due to, (4) 2
Nematologists, meeting, Puerto Rico, (2) 7
Netherlands, cattle export, (4) 21
fertiliser usage, (3) 41 BR
soil exhaustion in tree nurseries, (4) 2
New Zealand, fertiliser usage, (3) 41 BR
spraying crops by helicopter, (1) 25
hill pasture with weedkillers, (1) 25
Nigeria, antimalarial campaign, (1) 10
tsetse fly campaign, (4) 25
Nitra-Shell, aerial application, (2) 27
trial on ley, (3) 21 (fig.)
Nitrogenous fertilisers, (3) 42 BR
foliar application, (2) 2
for sisal, (1) 22
on grassland, U.K., (3) 21
Norway, fertiliser usage, (3) 41 BR
*Nosopsyllus* spp., (1) 13
Nozzle, design, in relation to drift, (4) 14
Nurseries, sisal, fertilisers for, (1) 22
soil exhaustion in tree plantings, (4) 2
Nut, hazel, value to Turkey, (1) 32
Nutrition research, tracer technique, (1) 5

## O

Oak, spraying against *Tortrix*, (3) 33
Oak leaf roller, (3) 33
Oats, grey speck disease, (4) 8
Oil-in-water emulsion, insecticidal, (3) 31
Oiled paper, as soil cover, (4) 5
Olive, production, Turkey, (1) 32
Olive fly, (2) 23
*Oncopeltus fasciatus*, (2) 29
Onions, weed control in, (4) 10
Orange, Mediterranean fruit fly on, (1) 18
Orchard, aerial spraying trials, (2) 27
Organophosphorus insecticides, (1) 3; (2) 33 BR
chlorinated terphenyl additives, (2) 13
in fly control, (1) 9
resistance to, in housefly, (1) 9; (2) 10
toxicology, (4) 39 BR
*See also* Malathion, Parathion, Phosdrin, Schradan.
*Orobanche ramosa*. *See* Broomrape.
*Ottelia alismoides*, control, (3) 27, 28 (fig.)
Overliming, boron deficiency and, (4) 7

## P

Paddy. *See* Rice.
Palm weevil, (2) 5
Pan American Sanitary Organisation, malaria eradication fund, (1) 27
Paraffinic oils, (3) 4
against banana leaf spot, (3) 2
Parathion, (2) 29, 30; (4) 39 BR
resistance to, in flies, (2) 10
resurgence of pests after, (3) 19
toxicity to coccinellids, (3) 18
Parsley, tolerance of urea spray, (2) 3
Pasture. *See* Grassland.
P.C.P., phytotoxicity, (1) 24
Pea, radiation mutant, (1) 4
weed control in, (4) 10, 11 (fig.)
Pea aphid, sensitive to Phosdrin, (2) 29
Pea curvature test, (4) 11 (fig.), 12
Peach, attack by Mediterranean fruit fly, (1) 17, 19 (fig.)
Pear, attack by Mediterranean fruit fly, (1) 17
*Pediculus humanus*, (1) 12
generations per year, (2) 11
Pentachlorophenol, (4) 10
toxicity to Salvinia, (3) 29
*Periplaneta americana*, (1) 13 (fig.)

*Persectania* spp., (1) 27
Perthane, (1) 13
Pest control, irradiation methods, (1) 6; (2) 13
research centres, (2) 33 BR; (3) 23
Pesticide, classification, U.S.A. (4) 35
synthesis and development of new, (3) 24
systemic, test, (3) 7
veterinary uses, (4) 39 BR
*See also* Fungicide, Insecticide, Nematocide, Weedkiller.
Pesticide residues, analysis, (2) 33 BR
bioassay, (2) 33 BR; (3) 24
estimation on crops, (1) 16 (fig.)
tolerance limits, (1) 15; (4) 34
toxicity to beneficial insects, (3) 18
Phenoxyacetic acid compounds, movement in plant tissues, (3) 5
Phosdrin, (2) 28
mammalian toxicity, (2) 30
residues, analysis, (3) 25 (fig.)
rapid disappearance, (1) 15
Phosphate fertilisers, (3) 42 BR
foliar application, (2) 2
Phosphorus ester insecticides. *See* Organophosphorus insecticides.
Phytosanitary regulations, (2) 22
Phytotoxicity, (4) 5
boron applications, (4) 7
insecticide solvents, (3) 31
mineral oils, (3) 2, 3
trace elements, (4) 6
Pick-up baler, (2) 18 (fig.) 19
Picking machines, for cotton, (3) 40 BR
*Pieris rapae*, (2) 30
Pineapple, foliar absorption of urea, (2) 4
Plant breeding, radiation-induced mutations, (1) 4
rice, (4) 28
sisal, (1) 24
Plant food ratio, change in, (3) 42 BR
Plant growth regulators, (4) 10
*See also* Weedkillers.
Plant protection, International co-operation, (2) 20
pest reporting service, (2) 23
Planting, interval after D-D fumigation, (4) 5
sisal, (1) 22
Plough, primitive type, (1) 31 (fig.)
Ploughing, methods in Turkey, (1) 31
wet rice, Indonesia, (4) 29 (fig.)
Plum, attack by Mediterranean fruit fly, (1) 17
*Plutella maculipennis*, (2) 30
Poisonous plants, (4) 39 BR
Poll strike of rams, (4) 31
Pollen, irradiation (1) 4
*Polygenis* sp., (1) 13
Pond weeds, control, (3) 26
Population pressure, correlated with fertiliser usage, (3) 41 BR
Portugal, oak spraying campaign, (3) 33
Potamogeton, control, (3) 27, 28
*Potamogeton nordosus*, (3) 29 (fig.)
Potash fertilisers, (3) 42 BR
foliar application, (2) 2, 4
Potato, aerial spraying, (2) 26 (fig.) 27
colorado beetle control, (2) 22
root eelworm, EPPO work, (2) 22
sprouting inhibited by irradiation, (1) 6
wart disease, EPPO work, (2) 22
Potato blight, control, (2) 27
Potato tissue block, (3) 7
*Pratylenchus penetrans*, (4) 3, 4
Pre-emergence treatment, in weed control, (4) 10
Pre-harvest drop retarders for, (4) 10
*Prodenia eridana*, (2) 29
Prolan, (2) 9
Propham, (4) 10
*Prunus avium*, (4) 3 (fig.) 4
*Prunus malaheb*, (4) 4
*Pseudaletia* sp., (1) 27
*Psoregates ovis*, (4) 32, 33 (fig.)
*Psorophora confinis*, (1) 11
*Pterocallidium* sp., (2) 29
*Pulex irritans*, (1) 13
*Pyrausta nubilalis*, (3) 32
Pyrethrins, (1) 9; (3) 13
Pyrethrum powder, (2) 13
toxicity to body louse, (1) 12
*Pyrus communis*, (4) 4

Quarantine regulations, citrus, (1) 18
Quince, attack by Mediterranean fruit fly, (1) 18

Radiation mutants, commercial growing, (1) 4
Radioactive isotope, agricultural uses, (1) 4
in pesticide research, (2) 33 BR
in weedkiller studies, (3) 6
Radioactive materials, supply of, (1) 5
Radioactivity, sterilization of insects by, (1) 6; (2) 13
Radiochemical Centre, Amersham, (1) 6 (fig.)
Raindrop, impact, (3) 35 (fig.) 37
Rainfall, soil erosion factor, (3) 34, 35
Rat cages, Wantage Irradiation laboratory, (1) 7
Reclamation disease, of cereals, (4) 7

Red Poll, cross with Brahman breed, (4) 21
Red ring disease, of coconut, (2) 5
Resistance to insecticides. *See* Insecticide resistance.
Resurgence, caused by spraying, definition, (3) 18
Rhodesia, soil conservation, (3) 34
*Rhopalopsylla claviculus*, (1) 13
*Rhyncophora palmarum*, (2) 5
Rice, (4) 27
  acreage, Turkey, (1) 32
  collecting transplanting material, (4) 27 (fig.)
  dieldrin dusting, (3) 32
  fertilizing, economics, (4) 30
  planting, Indonesia, (4) 28 (fig.)
  seed vernalisation, (4) 29
  selection and breeding, (4) 28
Rice stem borer, (4) 30
Root crops, boron deficiency, (4) 7
Root systems, in exhausted soil, (4) 2
Rooting of cuttings, stimulants, (4) 10
*Rosa canina*, (4) 4, 5 (fig.)
Rosetting, in apple, (4) 7 (fig.)
Rotation of crops, (4) 37 BR
Rush, eradication of tufted, (3) 26

## S

Sabi River Valley, catchment planning, (3) 38
*Salix* spp., (1) 26
*Salvinia auriculata*, (3) 26, 29
San José scale, (2) 22
Santa Gertrudis cattle, (4) 21
Scale insects, hydrogen-cyanide resistance, (2) 9
  ladybird predators of, (1) 20
Schradan, (3) 19
Screw worm, control, Curacao, (1) 6; (2) 13
*Scyphophorus acupunctatus*, (1) 22 (fig.) 23
Sealing light soils after D-D injection, (4) 5
Seed, radiation mutants, (1) 4
  vernalisation, (4) 29
Seed cotton, ginning, (3) 40 BR
Seed counter, (3) 24 (fig.)
Seed disinfectants, formulation, (3) 32
Selenium compounds, residues, zero tolerance, (4) 35
Sheep, ectoparasites, (4) 31
  statistics, Turkey, (1) 32
Sheep dips, (4) 32
  formulation, (3) 32
Shell A-A Weed Seed Killer, (2) 14
Shell Brush Killer, (1) 24
Shell Stand, Lisbon Conference, (4) 17 (fig.)
Shigellosis, in fly-infested villages, (1) 8
Shorthorn, cross with Brahman breed, (4) 21
Shamuti orange, (1) 18
Sieve tube, nature of, (3) 5
Sigatoka disease, of banana, (3) 2
  control, Hawaii, (2) 4
Sila-Master forage harvester, (2) 18 (fig.)
Silage, demonstration, Hatch Warren, (2) 16
Silage crops, harvesting, (2) 18
Silorator forage harvester, (2) 18 (fig.)
Simazin, (1) 3; (4) 26
Sisal, growing, East Africa, (1) 21
  response to nitrogen, (1) 22
  transplant, weevil damage, (1) 23 (fig.)
Sisal Research Station, Tanganyika, (1) 22, 23, 24
Sisal waste, manurial value, (1) 22
Sisal weevil, control, (1) 21, 22 (fig.) 23
Sleeping sickness, (4) 23, 25 (fig.)
Small holders, rice production, (4) 30
Smoke bomb, against tsetse fly, (4) 24
Sodium arsenite, (1) 27; (3) 5
Sodium chlorate, aerial application to pine plantings, New Zealand, (1) 27
Soil, biology, (2) 32 BR
  erosion losses, (3) 37
  fumigation, (4) 4
  temperature, D-D use and, (4) 5
Soil conservation, Rhodesia, (3) 34
Soil exhaustion, (4) 2, 3
Soil fertility, soil erosion and, (3) 36
Soil fungi, control, (2) 33 BR
Soil science, (2) 32, BR; 33 BR
Soil structure, rain drop and impact, (3) 37
*Solenopsis* sp., (2) 36
Solvent naphtha, in waterweed control, (3) 28
Solvents, emulsified, in weed control, (3) 28
  selection for insecticide formulation, (3) 31
Sowing, aerial, New Zealand, (1) 25
  interval after D-D fumigation, (4) 5
Southern army worm, (2) 29
Speckled yellows of sugar beet, (4) 8
Spotted lucern aphid, sensitive to Phosdrin, (2) 29
Spray drift. *See* Drift.
Spraying, (2) 33 BR
  aerial, (1) 25, 27; (2) 24, 36, (3) 33
    coverage test, (1) 26
  banana against leaf spot, (3) 2
  boom cover, (4) 16
  broomrape in tobacco, (2) 14
  field peas with DNBP, (1) 16 (fig.)
  hazards, (1) 14
  high- and low-volume compared, (2) 33 BR
  marsh Arab dwellings, (3) 43 BR
  nozzle function and design, (4) 14
  oaks, Portugal, (3) 33 (fig.)
  plants with trace elements, (2) 2
  sisal with dieldrin, (1) 23

Spraying machine, replaced by two-nozzle knapsack sprayer, (2) 34
Sayer exhaust sprayer, (2) 35
Sprouting, prevention by irradiation, (1) 6
Statistics, fertiliser production, (3) 41 BR
tea production, (4) 36 BR
Sterilisation, radiation, of food, (1) 6
of male insects, (1) 6; (2) 13
*Stethorus punctillum*, (3) 17 (fig.)
Stored product, pest control by irradiation, (1) 6
pests, EPPO report, (2) 23
Sudan, Gezira, weed control, (3) 27, 28
Sugar beet, (1) 32
boron deficiency, (4) 7
foliar absorption of nutrients, (2) 3
manganese deficiency, (4) 8
weedkillers for use in, (4) 10
Sulphur, wettable, compatible with urea in sprays, (2) 3
Sulphuric acid, weedkiller, (4) 9, 10
Superphosphate, aerial application, (1) 25
molybdenized, (4) 8
Swede, boron deficiency, (4) 7
Sweden, fertiliser usage, (3) 41 BR
Synergists, (1) 3; (2) 12
Systemic insecticides, (2) 33 BR; (3) 20

## T

2,4,5-T, aerial application, (1) 25 (fig.)
gorse eradication, New Zealand, (1) 26
toxicity to woody species, (1) 26
use in sisal field, (1) 24
Tachnid fly, (3) 17 (fig.)
Tanganyika, sisal growing, (1) 21
tsetse fly control, (4) 23
Tea, (4) 36 BR
blister blight, (2) 34
pests of, (4) 36 BR
production, Turkey, (1) 32
Temperature, of soil, D-D use and, (4) 5
viscosity of mineral oil sprays and, (3) 4
Tetraethyl pyrophosphate residues, zero tolerance, (4) 35
*Tetranychus* spp., resurgence, (3) 18
Thistle, eradication, (1) 25
Threshing, method, Turkey, (1) 31, 32 (fig.)
Ti-tree, eradication, (1) 26
*Tilapia melanopheura*, (3) 27
Tobacco, (1) 32
broomrape control in, (2) 14, 15 (fig.)
dieldrin dusting, (3) 32
erosion of ridges, (3) 37 (fig.)
seedbed alignment to lessen rain erosion, (3) 38 (fig.)
Tolerances, pesticide residues, (1) 15
Tomato, flower set, induced by growth regulators, (4) 10
foliar absorption of nutrients, (2) 3
Topdressing, (2) 24
grassland, (1) 25; (3) 21
*Tortrix viridana*, (3) 33
Toxaphene, resistance in housefly, (2) 10
toxicity to coccinellids, (3) 18
Toxicity, of agricultural chemicals, (1) 14
dieldrin emulsions in irrigation water, (3) 32
emulsified solvents to fish, (3) 28
Toxicity index, defined, (2) 29
Toxicology, veterinary, (4) 39 BR
Trace elements, aerial application, (2) 27
deficiency, (4) 6, 8
foliar absorption, (2) 4
Tracer technique, (1) 5
Tracklaying machines, stability, (1) 28
Tractor, (1) 31
stability, (1) 28, 29 (fig.)
Tractor driver, instruction, (1) 28
protection from toxic sprays, (1) 16 (fig.)
Transplanting, rice, (4) 28
Traps, insect, of plastics, (2) 36
Tree nurs ries, soil exhaustion, (4) 2
soil fumigation, (4) 4
Triazines, substituted, mode of action, (4) 26
Trichloroacetic acid, use in sisal nursery, (1) 24
in sugar beet seedbed, (4) 10
*Trichoplusia ni*, (2) 30
*Tridax procumbens*, (1) 24
Trinidad, antimalarial campaign, (1) 11
red ring of coconuts, (2) 5
*Trypanosoma brucei*, (4) 23
*Trypanosoma gambiense*, (4) 23
*Trypanosoma rhodesiense*, (4) 23
Trypanosomiasis, tsetse fly vector, (4) 23
Tristeza, of citrus, (2) 23
Trithion, (2) 29
Tropical agriculture, (4) 40 BR
Tropical Medicine, Lisbon Conference, (4) 17
Tsetse fly, control, (4) 24
distribution in Africa, (4) 23, 24
Turkey, agriculture in, (1) 30
Two-spotted spider mite, (2) 29

## U

*Ulex europaeus*, (1) 26
U.M.R., definition, (3) 2
*Unaspis euonymi*, (3) 18
United Kingdom, fertiliser usage, (3) 41 BR
silage demonstration, (2) 16

United States, Miller Amendment to Food and Drug Act, (4) 34
pesticide legislation, (1) 15
Unsulphonatable mineral residue, (3) 2
Urea, foliar application, (2) 2, 3
to tropical crops, (2) 4
granular, aerial application, (2) 27
movement in plant tissues, (3) 7

## V

*Vallisneria spiralis*, (3) 28 (fig.)
Vapam, (4) 4
Vegetables, foliar absorption of nutrients, (2) 3
fumigation with methyl bromide, (2) 36
Vernalisation, rice seed, (4) 29
Vigour tolerance, (2) 9; (3) 13
Virus diseases, of citrus, EPPO work, (2) 23
research at Rothamsted, (3) 15
Viscosity of mineral oils, in relation to sprayer, (3) 4

## W

Walnut, aerial spraying trial, (2) 27
Wantage, 'cobalt 60 field', (1) 5
Water buffalo, (3) 43 (fig.)
Water conservation, Rhodesia, (3) 36
Water-dispersible powders, (3) 31
Water hyacinth, spread of, (3) 29
Weather forecast, (4) 38 BR
Weed control, (3) 42 BR; (4) 9, 37 BR
in pine plantings, (1) 27
in rice field, (4) 30
in sisal nurseries, (1) 24
in waterways, (3) 26
plant structure as factor in, (4) 9
Weed cutting launch, (3) 27 (fig.)
Weedkillers, (1) 3; (3) 42 BR; (4) 9, 26
aerial application, (1) 25
application in fertiliser spray, (2) 3
assessment of selective, (4) 11 (fig.) 12
granular formulations, (4) 26
invert emulsions, (4) 26
mode of action, (2) 33 BR; (4) 12
movement in plant tissues, (3) 5
screening tests, (3) 24
specificity of butyric acid derivatives, (4) 12
systemic, (3) 7; (4) 10
toxicity to fish in ponds, (3) 28
*See also* 2,4-D, 2,4,5-T Dalapon, MCPA, MCPB, MCPP, Methoxone, Simazin, Sodium chlorate, Trichloroacetate.
Weeds, aquatic, (3) 26
eradication, (3) 27
flora, of Britain, (3) 42 BR
seed dormancy, (3) 29
*See also* Broomrape, Wild oats.
Weevils. *See* Coconut weevil; Sisal weevil.
West Germany, fertiliser usage, (3) 41 BR
Wheat, radiation-induced mutants, (1) 5
Wheat cylinder test, (4) 11 (fig.) 12
Whiptail, molybdenum deficiency sympton, (4) 8
Wild life, hazard from sprays doubtful, (1) 15
Wild oats, control in sugar beet, (4) 10
Willow, eradication, (1) 26
Wind effect, in spray drift, (4) 14, 16
World Health Assembly, 11th, (3) 44
World Health Organisation, concern with insecticide resistance, (1) 8; (2) 8
malaria eradication work reviewed, (3) 44
malaria fund, (1) 27
malaria world survey, (2) 4
rules for protection of spray operators, (1) 14
test for insecticide resistance of mosquitoes, (2) 13
Woodstock Agricultural Research Centre, (1) 16; (3) 22
Woody species, eradication by, 2, 4, 5-T (1) 26
*Wuchereria bancrofti*, (1) 12
*Wuchereria malayi*, (4) 18

## X

*Xenopsylla cheopsis*, (1) 13
*Xyleborus*, dieldrin spraying, (3) 18
Xyloporosis, of citrus, (2) 23

## Y

Yellow spot, molybdenum deficiency symptom, (4) 8
Yellow tip, of cereals, (4) 7
Yugoslavia, colorado beetle campaign, (2) 22

## Z

*Zebrina* test specimen for tracer studies, (3) 6
Zero tolerance, of pesticide residues, (4) 35
Zinc deficiency, (4) 8
apple, rosetting, (4) 7 (fig.)
Zinc sulphate, application methods, (4) 8
Zineb, (1) 3; (2) 3
Zuckerman working party on pesticide hazards, (1) 16
Zululand, tsetse fly campaign, (4) 24

# Span Index